

# perfect wine glass

Olaf Klinke \*

July 10, 2007

## 1 Introduction

The following is a description of the *perfect wine glass* variational problem. The objective is to find the surface which can enclose a maximal volume. What makes a possible solution different from the sphere is that we allow it to be *open* like a glass. I think one can restrict to surfaces of revolution, so the simplified version becomes:

**Problem 1.1** Let  $y : [0, \infty[ \rightarrow \mathfrak{R}$  be a continuous function with weak first derivative and compact support. The enclosed volume of the surface of revolution is (up to a constant)

$$V(y) = \int_0^\infty y(x)^2 dx$$

and the surface area is

$$A(y) = \int_0^\infty \sqrt{y(x)^2(1 + y'(x)^2)} dx$$

Now to get rid of the units take

$$v(y) = V(y)^{1/3}, a(y) = A(y)^{1/2}$$

and minimize the ratio

$$r(y) = a(y)/v(y). \tag{1}$$

Note that the functional  $r$  is invariant under scaling

$$y(x) \rightarrow c * y(x/c) \tag{2}$$

for positive real  $c$  and also invariant under  $y \rightarrow -y$ .

So one might want to consider the problem in a kind of projective space. I do not know whether minimizing  $r$  is equivalent to the problem

$$\min\{a(y) : v(y) = 1\} \tag{3}$$

but one could scale a minimizer for (1) as in (2) to obtain a function satisfying the constraint in (3).

To get rid of the condition of compact support we could alter the problem in the following way:

Consider pairs  $(y, x_0)$  where  $y$  is a real-valued weakly differentiable function on the positive real axis and  $x_0$  is some value  $> 0$ .

Then alter the functional  $A$  to

$$A(y, x_0) = \int_0^{x_0} \sqrt{y(x)^2(1 + y'(x)^2)} dx + 1/2 * y(x_0)^2,$$

---

\*Tulane University

which is thought of a surface of revolution with a flat bottom at  $x = x_0$ . Of course in the case  $y(x_0) = 0$  the new  $A$  is the same as the old one.

What might be a problem is that  $r$  is not defined for the zero function, and by the invariance (2) one can see that  $r$  attains any possible value in any neighborhood of zero, since a for a function  $y$  of bounded support

$$\lim_{c \rightarrow 0} c * y(x/c) = 0$$

for all  $x \geq 0$ , which means that we find a path where  $r$  is constant that links  $y$  and the zero function (at least in the topology of pointwise convergence).

I already did a simple discretization using B-splines and a gradient method, and my guess is that the half-sphere

$$y(x) = \sqrt{1-x}, x_0 = 1$$

is the global optimum.